

REFERÊNCIAS BIBLIOGRÁFICAS

FOLHETO APTAMIL SENSITIVE ACTIVE

1. HCP TRACKER,2019.
2. Morais. J Ped (Rio J).2016;92(Suppl 1):S46-S56.
3. Schmulson MJ, Drossman DA 2017.
4. Bellaiche M., et al. Acta Pædiatrica 2018 107, pp. 1276–1282
5. Bellaiche M., et al. Acta Paediatrica. 2020;00:1–8
6. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. Eur J Clin Nutr 2006, 60:1304-10.
7. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. J Pediatr Gastroenterol Nutr 2003, 36:343-51.
8. Havlicekova Z et al. Beta-palmitate - a natural component of human milk in supplemental milk formulas. Nutr J. 2016;15(1):28.
9. Litmanovitz I et al. Reduced crying in term infants fed high beta-palmitate formula: a double-blind randomized clinical trial. BMC Pediatr. 2014;14:152.
10. Yao M et al. Effects of term infant formulas containing high sn-2 palmitate with and without oligofructose on stool composition, stool characteristics, and bifidogenicity. J Pediatr Gastroenterol Nutr. 2014 ;59(4):440-8.
11. Nowacki J et al. Stool fatty acid soaps, stool consistency and gastrointestinal tolerance in term infants fed infant formulas containing high sn-2 palmitate with or without oligofructose: a double-blind, randomized clinical trial. Nutr J. 2014;13:105.
12. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. J Pediatr Gastroenterol Nutr. 2002;34(3):291-5
13. Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. J Pediatr Gastroenterol Nutr. 2011;52:238–225
14. Boehm G et al. Prebiotics in infant formulas. J Clin Gastroenterol. 2004; 38: S76-9.
15. Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. Br J Nutr. 2015;113(9):1339-44.
16. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. Am J Clin Nutr. 2013;98(2):561S-71S.
17. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. Nutr J 2007;6:8.
18. Savino F et al. Advances in the management of digestive problems during the first months of life. Acta Paediatr Suppl 2005, 94:120.
19. Savino F et al. “Minor” feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. Acta Paediatr Suppl 2003, 91:86-90.

20. Salvatore S et al Review shows that parental reassurance and nutritional advice help to optimise the management of functional gastrointestinal disorders in infants Acta Pædiatrica 2018 107 pp 1512 1520.
21. Comparativo de produtos de mesma categoria realizado em Junho de 2024.
22. Vandenplas Y et al Functional gastrointestinal disorders in infancy impact on the health of the infant and Family Pediatr Gastroenterol Hepatol Nutr 2019.
23. Gordon M, et al. Dietary modifications for infantile colic. Cochrane Database of Systematic Reviews 2018, Issue 10. Art. No.: CD011029.
24. Meyer, R. et al. Systematic review of the impact of feed protein type and degree of hydrolysis on gastric emptying in children. BMC Gastroenterol. 2015
25. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. Acta Pædiatrica 2015.
26. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. Ital J Gastroenterol Hepatol, 1997; 29:237-42.
27. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. Journal of Pediatrics, 2003;111(4):355-359.
28. Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. Pediatr. Mod. 2014; 50(8): 339-342.
29. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. J Med Assoc Thai. 2004 Jun;87(6):641- 9.