

REFERÊNCIAS BIBLIOGRÁFICAS

Referências – Material extensamente hidrolisado_CL06

1. Vandenplas Y, Broekaert I, Domellöf M, Indrio F, Lapillonne A, Pienar C, et al. An ESPGHAN position paper on the diagnosis, management and prevention of cow's milk allergy. *Journal of Pediatric Gastroenterology & Nutrition*. 2023.
2. Estudo observacional, prospectivo, aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil. 2016: dados preliminares.
3. Epifanio, M. et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. *Biomed J Sci & Tech Res* 25(1)-2020. BJSTR. MS.ID.00416.
4. Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *J Biol Regul Homeost Agents*. 2012;26(3 Suppl):49-59.
5. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr*. 2008;138(6):1091-5.
6. Moro G, Arslanoglu S, Stahl B, Jelinek J, Wahn U, 12. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child*. 2006;91(10):814-9.
7. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr*. 2005;41(5):584-99. Leia menos
8. Ballard O, Morrow AL. *Pediatr Clin North Am*. 2013;60(1):49-74.
9. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
10. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics*. 2013;131(1):262-72.
11. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
12. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients*. 2021; 13: 247.
13. Marten B et al. Medium-chain triglycerides. *International Dairy Journal*. 2006;16: 1374-1382
14. Solé D et al. Consenso brasileiro sobre alergia alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol*. 2018;2(1):7-82.
15. Venter C et al. Diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: a UK primary care practical guide. *Clin Transl Allergy*. 2013;3(1):23.
16. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.

17. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol.* 2012;23(4):307–14.
18. Joosten K, Meyer R. Nutritional screening and guidelines for managing the child with faltering growth. *Eur J Clin Nutr.* 2010;64:S22–4.
19. Joint WHO/FAO/UNU Expert Consultation. Protein and amino acid requirements in human nutrition: report of a joint. WHO Tech Rep Ser [Internet]. 2007;1–265. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18330140>
20. Lucia Diaferio L et al. May Failure to Thrive in Infants Be a Clinical Marker for the Early Diagnosis of Cow's Milk Allergy?. *Nutrients.* 2020 Feb 13;12(2):466. doi: 10.3390/nu12020466.
21. Boaventura RM, Mendonça RB, Fonseca FA, Mallozi M, Souza FS, Sarni ROS. Nutritional status and food intake of children with cow's milk allergy. *Allergol Immunopathol (Madr).* 2019;47(6):544-550. doi:10.1016/j.aller.2019.03.003
22. Eveleens, R D et al. Definitions, predictors, and outcomes of feeding intolerance in critically ill children: A systematic review. *Clin Nutr,* 2020, 39.3: 685-693.
23. Marino, LV et al. Feeding intolerance in children with critical illness. *Clin. Nutr.,* 2020, 39.3: 609-611.
24. IQVIA outubro de 2023.
25. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr.* 1995;49 Suppl 1:S39 S48.
26. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol.* 2001;12(2):83 86.
27. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics,* 90(3), 463 465.
28. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics,* 127(4), 550 557.